

REVIEW

Should we be testing for urogenital *Mycoplasma hominis*, *Ureaplasma parvum* and *U. urealyticum* in men and women? – a Position Statement from the European STI Guidelines Editorial Board

P. Horner,^{1,2} G. Donders,³ M. Cusini,⁴ M. Gomberg,⁵ J.S. Jensen,⁶ M. Unemo^{7,*}

¹Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, United Kingdom

²National Institute for Health Research Health Protection Research Unit in Evaluation of Interventions, University of Bristol, Bristol, United Kingdom

³Department of Obstetrics and Gynecology, University Hospital Antwerp, Edegem, Belgium

⁴Department of Dermatology, Fondazione IRCCS Ca' Granda Ospedale Policlinico, Milano, Italy

⁵Moscow Scientific and Practical Center of Dermatovenereology and Cosmetology, Moscow, Russia

⁶Infection Preparedness, Research Unit for Reproductive Tract Microbiology, Statens Serum Institut, Copenhagen, Denmark

⁷World Health Organization Collaborating Centre for Gonorrhoea and Other STIs, Department of Laboratory Medicine, Microbiology, Faculty of Medicine and Health, Örebro University, Sweden

*Correspondence to: M Unemo. E-mail: magnus.unemo@regionorebrolan.se

Magnus Unemo, World Health Organization Collaborating Centre for Gonorrhoea and Other

STIs, Department of Laboratory Medicine, Microbiology, Örebro University Hospital, SE-701 85 Örebro, Sweden. Tel.: +46 19 6022038. Fax: +46 19 127416

Running head: Testing for urogenital mycoplasmas in men and women?

Word count: 267 words (abstract), 2916 words (manuscript); **Tables:** 1; **Figures:** 0

Conflicts of interest

The authors have no conflicts of interest related to this paper.

Funding sources

None.

Abstract

At present, we have no evidence that we are doing more good than harm detecting and subsequently treating *Mycoplasma hominis*, *Ureaplasma parvum* and *Ureaplasma urealyticum* colonisations/infections. Consequently, routine testing and treatment of asymptomatic or symptomatic men and women for *M. hominis*, *U. urealyticum*, and *U. parvum* is not recommended. Asymptomatic carriage of these bacteria is common and the majority of individuals do not develop disease. Although *U. urealyticum* has been associated with urethritis in men, it is probably not causal unless a high load is present (likely carriage in 40-80% of detected cases). The extensive testing, detection and subsequent antimicrobial treatment of these bacteria performed in some settings may result in selection of antimicrobial resistance, in these bacteria, “true” STI agents, as well as in the general microbiota, and substantial economic cost for society and individuals, particularly women. The commercialisation of many PCR assays detecting traditional non-viral STIs together with *M. hominis*, *U. parvum* and/or *U. urealyticum* have worsened this situation. Thus, routine screening of asymptomatic men and women or routine testing of symptomatic individuals for *M. hominis*, *U. urealyticum*, and *U. parvum* is not recommended. If testing of men with symptomatic urethritis is undertaken, traditional STI urethritis agents such as *Neisseria gonorrhoeae*, *Chlamydia trachomatis*, *M. genitalium* and, in settings where relevant, *Trichomonas vaginalis* should be excluded prior to *U. urealyticum* testing and quantitative species-specific molecular diagnostic tests should be used. Only men with high *U. urealyticum* load should be considered for treatment, however, appropriate evidence for effective treatment regimens is lacking. In symptomatic women, bacterial vaginosis (BV) should always be tested for and treated if detected.

Key messages

- Routine screening of asymptomatic men and women or routine testing of symptomatic individuals for *Mycoplasma hominis*, *Ureaplasma urealyticum*, and *Ureaplasma parvum* is not recommended.
- The extensive testing, detection and antimicrobial treatment of urogenital *M. hominis*, *U. parvum* and *U. urealyticum* performed in some settings result in a substantial burden and economic cost for society and individuals, particularly women. Instead, the diagnostics and treatment of traditional, more severe “true” STIs and BV in symptomatic women need to be improved.
- *U. urealyticum* in high bacterial loads might cause a small proportion of male NGU, but the majority of men and women infected/colonised with *U. urealyticum* do not develop disease. Antimicrobial treatment resulting in eradication is difficult, eradication is not strongly associated with cure, and treatment may select/induce resistance in urogenital mycoplasmas and other bacteria including the more severe “true” STI agents.

Introduction

Mycoplasmas and ureaplasmas belong to the class Mollicutes. *Mycoplasma genitalium* is a “true” STI causing male urethritis, and is associated with cervicitis and an increased risk of pelvic inflammatory disease (PID), endometritis and infertility.^{1,2} However, *Mycoplasma hominis*, *Ureaplasma urealyticum* (previously *U. urealyticum* biovar 2) and *U. parvum* (earlier *U. urealyticum* biovar 1)³ are frequently found in the human urogenital tract in both healthy individuals and symptomatic patients.⁴ In several settings, comprehensive testing and subsequent antimicrobial treatment of these three urogenital mycoplasma species in adults is performed. In many countries, this testing has also increased due the introduction of multiplex

PCR assays detecting traditional non-viral “true” STI agents together with *M. hominis*, *U. parvum* and/or *U. urealyticum*.⁵⁻⁷ Nevertheless, the evidence base for these three mycoplasmas as aetiological agents of STI syndromes and complications in adult men and women can be questioned. Most older studies used culture and this is still commonly used due to the availability of simple and easy to use culture kits with inappropriate antimicrobial susceptibility testing. However, culture does not distinguish between *U. urealyticum* and *U. parvum*, and results are often reported as *U. urealyticum* instead of *Ureaplasma* spp. leading to further confusion. Qualitative PCR assays are also commonly used without species differentiation and with inappropriate reporting. Furthermore, in most studies, the strong association between bacterial vaginosis (BV) in “patient” and/or sexual partner has not been adjusted for. This is particularly an issue for *M. hominis* but also for ureaplasmas.^{4,8-10} These and additional confounding factors make interpretation of many previous studies exceedingly difficult.

We reviewed the evidence for *M. hominis*, *U. parvum* and *U. urealyticum* as aetiological agents of urethritis, cervicitis and additional STI syndromes and complications in adult men and non-pregnant women, and conclude when these species should be considered for testing. We also suggest further research essential to provide appropriate evidence for unresolved questions. To avoid some of the confounding factors, we focused on international peer-reviewed papers using molecular diagnostics and appropriate species differentiation. Relevance of these bacteria in pregnancy or in neonates was not addressed, because this has been reviewed recently elsewhere.¹¹⁻¹⁵

Men

Male urethritis

There is no evidence from case control studies that *M. hominis* causes non-gonococcal urethritis (NGU).¹⁶⁻²⁰ It appears to be a relatively uncommon microorganism in men attending departments of sexual health (2-4%), although colonisation can be as high as 20%.^{16-18,21}

U. urealyticum and *U. parvum* can both be detected in men with and without NGU. Earlier studies did not differentiate between *U. urealyticum* and *U. parvum*, which continues to be the case if culture alone is used.^{9,10} *U. parvum* is detected more often in controls than cases in most studies, which probably explains why earlier studies failed to demonstrate a consistent association of ureaplasmas with NGU.^{9,10}

The population prevalence of *U. parvum* in men is unknown but it is likely more common than *U. urealyticum* as it is detected more frequently in men without urethritis than *U. urealyticum*.^{9,22} A recent meta-analysis of case control studies demonstrated no association of *U. parvum* with NGU.⁹ This was also observed by Frølund et al,²² but not in a few other studies of non-gonococcal non-chlamydial urethritis where *U. parvum* was associated with disease, in particular when present in high loads.^{21,23,24} Additional large and well-designed studies using quantitative molecular detection of *U. parvum* with appropriate cut-off for high bacterial load in men with symptomatic urethritis might be valuable.

The population prevalence of *U. urealyticum* is unknown but is probably 5-15% in men aged 16-44 years old,^{21,22,25,26} being more common in younger men and associated with a recent change in sexual partner.^{26,27} *U. urealyticum* is associated with NGU. However, although detected in 5-24% of men with NGU it is probably only causal in 3-11% of NGU cases, i.e. in 40-80% of cases it is probably only carriage.^{9,21,22,25,28,29} A recent meta-analysis demonstrated a significant association with 18.3% of men with NGU and 13.7% of controls being *U. urealyticum*-positive with a pooled odds ratio (OR) of 1.57 (95% CI: 1.05–2.35), $p=0.029$.⁹ Although, NGU caused by *U. urealyticum* is more likely to develop in younger men, the majority of men carrying *U. urealyticum* will not develop NGU. The development of

NGU is associated with a higher bacterial load and fewer lifetime sexual partners.^{22-24,30,31} As *U. urealyticum* carriage in men without urethritis is associated with younger age,^{26,27} this suggests that the adaptive immunity attenuates the clinical manifestation of *U. urealyticum* infection; repeated or prolonged exposure to *U. urealyticum* via multiple sex partners may result in either asymptomatic colonisation without signs of urethral inflammation or shorter duration of symptoms.^{22,27,31} Using quantitative molecular detection of *U. urealyticum* with appropriate cut-off for high bacterial load in men with symptomatic urethritis can significantly increase the positive predictive value.^{22,23,30,32} However, additional studies using different quantitative molecular tests and examining symptomatic and asymptomatic male populations in different settings are required before any exact cut-off levels can be recommended.

Male infertility

A recent meta-analysis³³ as well as two studies (which did not exclude “true” STIs or BV, and only included *M. hominis* culture positive samples)^{34,35} have suggested an association of *M. hominis* with infertility in men. However, *M. hominis* is strongly associated with several “true” STIs that can cause infertility as well as with BV,^{4,36} which is common in women, and two recent studies indicated that sexual partners share their genital tract microbiome, suggesting that molecular detection in men is likely to reflect the carriage in their female sexual partner.^{37,38} BV is more common in women with infertility and is associated with tubal factor infertility as well as with poor implantation of the embryo as suggested by a study of women undergoing in vitro fertilisation (IVF).^{39,40} Thus, considerable caution should be exercised in attributing the detection of *M. hominis* as causal of male infertility before additional studies have been performed. These studies should be appropriately designed and use quantitative PCR, and address “true” STIs and BV as confounders (in infertile men and

partners) as well as showing that treating the *M. hominis* infection in infertile men will restore fertility.

A recent meta-analysis demonstrated no association with *U. parvum* but suggested an association between *U. urealyticum* and male infertility.³³ Of the five included studies, three were from China where a high prevalence was observed in both cases (19.6%) and controls (8.3%)³³ compared to a study from Jordan 1.1% vs 2.9%⁴¹ and Iran 9% vs 1%,⁴² respectively. Whether *U. urealyticum* actually causes male infertility remains unclear, some studies do not differentiate *U. urealyticum* and *U. parvum*, further complicating interpretation of the data.^{34,43-45} Possible explanations for an inconsistent association in case control studies of male infertility include, failure to differentiate *U. urealyticum* and *U. parvum*^{44,45} and association by confounding as *U. urealyticum* is associated with younger age, recent change in sexual partner and fewer lifetime sexual partners and the association of ureaplasmas with BV.^{4,26,27,39}

Women

The prevalence of *M. hominis*, *U. urealyticum* and *U. parvum* in non-pregnant sexually active symptomatic and asymptomatic women, measured by molecular tests including species differentiation, has ranged between 3.1-15%, 5.2-20%, and 20-89%, respectively.^{5,46-54} The large variation in prevalence probably reflect both methodological and true population differences, in particular in the prevalence of BV, the most important confounder. *M. hominis* and ureaplasmas can be horizontally transmitted and although colonization tends to decrease with age until puberty, detection of these bacteria in prepubertal girls even in the absence of sexual abuse is not unusual,^{55,56} which illustrates that sexual transmission is not required. Nevertheless, among adults most cases of new colonisation with *M. hominis* and ureaplasmas occurs from sexual contact⁵⁷ and is correlated to the number of sexual partners.⁵⁸

Overall, in symptomatic women with dysuria, vaginal discharge, painful intercourse, and/or lower abdominal pain the spectrum of symptoms do not differ in ureaplasma-negative women compared to women positive for *U. urealyticum* or *U. parvum*.⁴⁸ However, both of these ureaplasmas are frequently associated with increased positivity for several traditional STIs, e.g. *C. trachomatis* and *M. genitalium*, and/or BV.^{8,46,49,59} The bacterial load of particularly *M. hominis* and to a lesser extent *U. parvum* and *U. urealyticum* can be significantly increased in the dysbiosis of BV.^{4,48,60} However, despite the association between particularly *M. hominis* and BV, *M. hominis* cannot be detected in approximately one third of women with BV and, accordingly, it is neither a sufficiently sensitive nor specific bacterial marker for diagnosis of BV.^{8,61-63} Despite not being susceptible to metronidazole, eradication or a decrease in the *M. hominis* load after BV treatment has also been reported,⁶⁴⁻⁶⁶ further indicating that *M. hominis* frequently belongs to the dysbiosis of BV. BV treatment studies using quantitative molecular detection methods for *M. hominis*, *U. urealyticum* and *U. parvum* are required. In many studies, appropriate species differentiation of *U. urealyticum* and *U. parvum* have not been performed and/or traditional STIs and especially BV have not been addressed as confounding factors, making disease association with the urogenital mycoplasmas exceedingly difficult.

Vulvovaginitis

There are no case control studies or other appropriate evidence that *M. hominis*, *U. parvum* or *U. urealyticum* causes an inflammatory vulvovaginitis.^{4,48,50} The number of leukocytes in vaginal smears are also not increased in women positive for only ureaplasmas.⁴⁸

Cervicitis

No case control studies using sensitive and specific molecular diagnostic tests have provided appropriate evidence that *M. hominis*, *U. parvum* or *U. urealyticum* causes cervicitis. For example, the unadjusted prevalence ratios of cervicitis have been reported as 1.00, 1.09, and 0.96 for *M. hominis*, *U. parvum* or *U. urealyticum*, respectively.⁶⁷ Also in additional cervicitis studies, none of these three urogenital mycoplasmas was associated with cervicitis⁶⁸ and the bacterial load of neither *U. parvum* nor *U. urealyticum* has been associated with symptoms or signs of genital infection.⁴⁹ Nevertheless, in one molecular study of non-gonococcal non-chlamydial cervicitis, despite no difference in *U. parvum* and *U. urealyticum* presence in women with cervicitis and controls⁶⁹ the bacterial load of *U. parvum* and *U. urealyticum* were significantly higher in women with cervicitis compared to controls.⁶⁹

Female urethritis and urethral pain syndrome

Appropriate studies are mainly lacking, however, no case control or other studies providing evidence that *M. hominis*, *U. parvum* or *U. urealyticum* causes urethritis in women are available. One study of the urethral pain syndrome in women showed that 46% of women with urethral pain carried *Ureaplasma* species compared with 64% of the controls. The prevalence of *U. parvum* and *U. urealyticum* were similar in women with the urethral pain syndrome and controls.⁵¹ Using undifferentiated quantitative ureaplasma culture, early work suggested some evidence of a role of high bacterial loads in women with acute urethral syndrome.⁷⁰ Studies using up-to-date quantitative techniques for ureaplasma detection are recommended.

Pelvic Inflammatory Disease (PID), salpingitis and infertility

Studies are few and no case control studies have yet provided appropriate evidence that *M. hominis*, *U. parvum* or *U. urealyticum* cause PID, salpingitis or infertility.^{71,72} Although *M. hominis* has been isolated from laparoscopically obtained samples, it was always found also in

the vagina, so it may well be present in a background of BV associated bacteria which were not cultured.^{73,74} In another study, the detection of *M. hominis* in the lower genital tract was not associated with *C. trachomatis*-negative and gonorrhoea-negative salpingitis and was not isolated from the salpinges indicating that it is unlikely to be causal.⁷¹ However, it is occasionally the sole pathogen isolated from the upper genital tract.⁷⁴ In infertility, pooled data for non-pregnant women were analysed in a systematic review,⁵² and both *M. hominis* (11.5% vs. 14.5%, $p=0.03$) and *U. urealyticum* (19.5% vs. 25.0%, $p=0.004$) were more common among asymptomatic women presenting for infertility ($n=1205$) compared with symptomatic women ($n=1131$; with vulvovaginitis signs), possibly indicating an association with infertility. In general, *C. trachomatis* infection, gonorrhoea and/or BV as confounding factors have been present or not appropriately excluded in most studies, and BV is strongly associated with infertility.⁴⁰ Microbiota studies of invasive samples in women with verified PID, e.g. laparoscopically taken specimens, would be valuable to adequately address this as the BV associated bacteria are often uncultivable.

Ectopic pregnancy

There is no clear evidence that any of the urogenital mycoplasmas, including the “true” STI agent *M. genitalium*, result in ectopic pregnancy.⁷⁵

Discussion and conclusions

In men, *M. hominis* does not cause disease and is probably mostly a reflection of BV in the partner and the presence of *U. parvum* is not evidently associated with NGU or infertility.^{9,16-18,22} *U. urealyticum* is associated with a small proportion of NGU cases, in particular in younger men with fewer lifetime sexual partners and a high *U. urealyticum* load. However, in

~40-80% of cases where it is detected, it is not the aetiological agent.^{9,21-23,25,28-32} It remains unlikely that *U. urealyticum* can cause infertility.

In women, there is no adequate evidence that *M. hominis*, *U. parvum* or *U. urealyticum* causes an inflammatory vulvovaginitis, cervicitis, urethritis, PID or infertility.^{4,48-51,67-69,71,72,76,77} In many studies, appropriate species differentiation of *U. urealyticum* and *U. parvum* has not been performed and/or important confounding factors such as recognized STIs and especially BV have not been addressed, making disease associations with the urogenital mycoplasmas mostly undocumented.

There are no international evidence-based management guidelines for *M. hominis*, *U. parvum* and *U. urealyticum*, and appropriate evidence for effective treatment regimens is lacking. Because mycoplasmas lack the rigid cell wall of other bacteria, they are intrinsically resistant to β -lactam antimicrobials, such as penicillins and cephalosporins, and other antimicrobials targeting the cell wall. *M. hominis* is additionally naturally resistant to 14- and 15-membered macrolides (azithromycin, clarithromycin, erythromycin), but not to 16-membered macrolides such as josamycin and the *in vitro* susceptibility to doxycycline is high for strains lacking the *tetM* gene. *U. urealyticum* is moderately sensitive to 14-membered macrolides. In general, urogenital *M. hominis*, *U. parvum* and *U. urealyticum* can be difficult to eradicate in many individuals because of true antimicrobial resistance but also because of lower activity of the antimicrobials at low pH and lack of bactericidal activity.^{4,27,78-80} Additionally, suboptimal antimicrobial susceptibility testing methods, including many commercial kits, are frequently used.⁸⁰ The extensive treatment of these commonly colonising commensals with suboptimal antimicrobial regimens selects for antimicrobial resistance in these bacteria and in many of the more severe bacterial “true” STI agents as well as in the general microbiota. Overall, the extensive testing, detection and subsequent antimicrobial treatment of urogenital *M. hominis*, *U. parvum* and *U. urealyticum* in some settings result in a

substantial burden and economic cost for society (e.g. unnecessary use of diagnostic tests, health care visits, antimicrobial misuse, and emergence of antimicrobial resistance) and individuals (e.g. economical burden, stigmatization, anxiety, and possibly breakdown of relationships including marriages). The commercialisation of many PCR assays detecting traditional non-viral STIs together with *M. hominis*, *U. parvum* and/or *U. urealyticum* have worsened this situation. At present, we have no evidence that we are doing more good than harm detecting and subsequently treating these bacteria.

Should testing for *M. hominis*, *U. urealyticum*, and *U. parvum* be undertaken in STI syndromes?

- *U. urealyticum* in high bacterial loads might cause a small proportion of male NGU, but the majority of men and women infected/colonised with *U. urealyticum* do not develop disease. Antimicrobial treatment which results in eradication is difficult^{4,27,78,79} and cure is not associated with eradication.⁷⁸ Treatment may result in development of antimicrobial resistance in urogenital mycoplasmas but also in other bacteria including the traditional, more severe “true” STI agents. Routine testing and/or treatment is therefore not recommended. If testing of men with symptomatic urethritis is undertaken, traditional STI urethritis agents such as *N. gonorrhoeae*, *C. trachomatis*, *M. genitalium* and, in settings where relevant, *Trichomonas vaginalis* should be excluded prior to *U. urealyticum* testing and quantitative molecular diagnostic tests should be used. Only men with high *U. urealyticum* load should be considered for treatment, however, appropriate evidence for effective treatment regimens is lacking.

- Testing for *M. hominis* and *U. parvum* and subsequent antimicrobial treatment of positive men or women is currently not recommended. Instead, “true” STIs and BV in symptomatic women should be diagnosed and treated.

Well-designed, large, randomised controlled studies to investigate unresolved issues regarding *M. hominis*, *U. parvum* and/or *U. urealyticum* and their independent associations with STI syndromes and complications such as possibly infertility,³³⁻³⁵ PID and prostate cancer⁸¹⁻⁸⁴ could be valuable. In these studies, it is recommended to control age, sexual behaviour (number and change of sexual partners), use quantitative molecular diagnostic tests investigating bacterial load, distinguish *U. urealyticum* and *U. parvum*, and exclude traditional STIs such as gonorrhoea, chlamydia, *M. genitalium* and trichomoniasis. Furthermore, it is crucial to address aerobic vaginitis and particularly BV and ideally also the specific BV-associated bacteria in controls and symptomatic individuals positive for urogenital mycoplasmas and their sexual partners. It is also important to show that antimicrobial treatment eradicates the mycoplasmas and that lack of eradication is associated with persistent symptoms and signs, documenting that it is not only an effect of treating a general dysbiosis.

Acknowledgements

We are grateful for valuable input on the present position statement to Keith Radcliffe, Jonathan Ross, Allesandra Sensini, Agnieszka Serwin, and Andy Winter.

Current composition of the European Guideline Editorial Board can be found at http://www.iusti.org/regions/Europe/pdf/2013/Editorial_Board.pdf.

List of contributing organizations and membership can be reviewed at www.iusti.org/regions/Europe/euroguidelines.htm.

PH is supported by the National Institute for Health Research Health Protection Research Unit (NIHR HPRU) in Evaluation of Interventions at the University of Bristol in partnership with Public Health England (PHE). The laboratory of MU is supported by the WHO, ECDC, Örebro County Council Research Committee, Örebro, Sweden, and Foundation for Medical Research at Örebro University Hospital, Örebro, Sweden. However, no funding was received for the present Position Statement and the views expressed are only those of the authors.

References

1. Unemo M, Jensen JS. Antimicrobial-resistant sexually transmitted infections: gonorrhoea and *Mycoplasma genitalium*. *Nat Rev Urol* 2017; **14**: 139-152.
2. Jensen JS, Cusini M, Gomberg M, Moi H. 2016 European guideline on *Mycoplasma genitalium* infections. *J Eur Acad Dermatol Venereol* 2016; **30**: 1650-1656.
3. Robertson JA, Stemke GW, Davis JW, Jr. *et al.* Proposal of *Ureaplasma parvum* sp. nov. and emended description of *Ureaplasma urealyticum* (Shepard *et al.* 1974) Robertson *et al.* 2001. *Int J Syst Evol Microbiol* 2002; **52**: 587-597.
4. Taylor-Robinson D. Mollicutes in vaginal microbiology: *Mycoplasma hominis*, *Ureaplasma urealyticum*, *Ureaplasma parvum* and *Mycoplasma genitalium*. *Res Microbiol* 2017; **168**: 875-881.
5. Kim Y, Kim J, Lee KA. Prevalence of sexually transmitted infections among healthy Korean women: implications of multiplex PCR pathogen detection on antibiotic therapy. *J Infect Chemother* 2014; **20**: 74-76.
6. Fernández G, Martró E, González V *et al.* Usefulness of a novel multiplex real-time PCR assay for the diagnosis of sexually-transmitted infections. *Enferm Infecc Microbiol Clin* 2016; **34**: 471-476.

7. Del Prete R, Ronga L, Lestingi M *et al.* Simultaneous detection and identification of STI pathogens by multiplex real-time PCR in genital tract specimens in a selected area of Apulia, a region of Southern Italy. *Infection* 2017; **45**: 469-477.
8. Malaguti N, Bahls LD, Uchimura NS, Gimenes F, Consolaro ME. Sensitive detection of thirteen bacterial vaginosis-associated agents using multiplex polymerase chain reaction. *Biomed Res Int* 2015; **2015**: 645853.
9. Zhang N, Wang R, Li X, Liu X, Tang Z, Liu Y. Are *Ureaplasma* spp. a cause of nongonococcal urethritis? A systematic review and meta-analysis. *PLoS One* 2014; **9**: e113771.
10. Horner PJ, Blee K, Falk L, van der Meijden W, Moi H. 2016 European guideline on the management of non-gonococcal urethritis. *Int J STD AIDS* 2016; **27**: 928-937.
11. Taylor-Robinson D, Lamont RF. Mycoplasmas in pregnancy. *BJOG* 2011; **118**: 164-174.
12. Donders GGG, Ruban K, Bellen G, Petricevic L. Mycoplasma/Ureaplasma infection in pregnancy: to screen or not to screen. *J Perinat Med* 2017; **45**: 505-515.
13. Sweeney EL, Dando SJ, Kallapur SG, Knox CL. The human ureaplasma species as causative agents of chorioamnionitis. *Clin Microbiol Rev* 2016; **30**:349-379.
14. Viscardi RM. *Ureaplasma* species: role in neonatal morbidities and outcomes. *Arch Dis Child Fetal Neonatal Ed* 2014; **99**:F87-92.
15. Viscardi RM, Kallapur SG. Role of *Ureaplasma* respiratory tract colonization in bronchopulmonary dysplasia pathogenesis: Current concepts and update. *Clin Perinatol* 2015; **42**:719-738.
16. Bowie WR, Pollock HM, Forsyth PS *et al.* Bacteriology of the urethra in normal men and men with nongonococcal urethritis. *J Clin Microbiol* 1977; **6**: 482-488.

17. Holmes KK, Handsfield HH, Wang SP *et al.* Etiology of nongonococcal urethritis. *New Engl J Med* 1975; **292**: 1199-1205.
18. Deguchi T, Yoshida T, Miyazawa T *et al.* Association of *Ureaplasma urealyticum* (biovar 2) with nongonococcal urethritis. *Sex Transm Dis* 2004; **31**: 192-195.
19. Bachmann LH, Manhart LE, Martin DH *et al.* Advances in the understanding and treatment of male urethritis. *Clin Infect Dis* 2015; **61 Suppl 8**: S763-769.
20. Totten PA, Taylor-Robinson D, Jensen JS. Genital mycoplasmas. In: Holmes KK *et al.*, eds. Sexually transmitted diseases, 4th ed. New York, McGraw-Hill Medical, 2008:709–736.
21. Cox C, McKenna JP, Watt AP, Coyle PV. *Ureaplasma parvum* and *Mycoplasma genitalium* are found to be significantly associated with microscopy-confirmed urethritis in a routine genitourinary medicine setting. *Int J STD AIDS* 2016; **27**: 861-867.
22. Frølund M, Lidbrink P, Wikström A, Cowan S, Ahrens P, Skov Jensen J. Urethritis-associated pathogens in urine from men with non-gonococcal urethritis: A case-control study. *Acta Derm Venereol* 2016; **96**: 689-694.
23. Strauss M, Colodner R, Sagas D, Adawi A, Edelstein H, Chazan B. Detection of *Ureaplasma* species by a semi-quantitative PCR test in urine samples: Can it predict clinical significance? *Isr Med Assoc J* 2018; **1**: 9-13.
24. Deguchi T, Shimada Y, Horie K *et al.* Bacterial loads of *Ureaplasma parvum* contribute to the development of inflammatory responses in the male urethra. *Int J STD AIDS* 2015; **26**: 1035-1039.
25. Khatib N, Bradbury C, Chalker V *et al.* Prevalence of *Trichomonas vaginalis*, *Mycoplasma genitalium* and *Ureaplasma urealyticum* in men with urethritis attending an urban sexual health clinic. *Int J STD AIDS* 2015; **26**: 388-392.

26. Jensen AJ, Kleveland CR, Moghaddam A, Haaheim H, Hjelmevoll SO, Skogen V. *Chlamydia trachomatis*, *Mycoplasma genitalium* and *Ureaplasma urealyticum* among students in northern Norway. *J Eur Acad Dermatol Venereology* 2013; **27**: e91-e96.
27. Horner P, Thomas B, Gilroy CB, Egger M, Taylor-Robinson D. Role of *Mycoplasma genitalium* and *Ureaplasma urealyticum* in acute and chronic nongonococcal urethritis. *Clin Infect Dis* 2001; **32**: 995-1003.
28. Couldwell DL, Gidding HF, Freedman EV *et al.* *Ureaplasma urealyticum* is significantly associated with non-gonococcal urethritis in heterosexual Sydney men. *Int J STD AIDS* 2010; **21**: 337-341.
29. Yoshida T, Ishiko H, Yasuda M *et al.* Polymerase chain reaction-based subtyping of *Ureaplasma parvum* and *Ureaplasma urealyticum* in first-pass urine samples from men with or without urethritis. *Sex Transm Dis* 2005; **32**: 454-457.
30. Shimada Y, Ito S, Mizutani K *et al.* Bacterial loads of *Ureaplasma urealyticum* contribute to development of urethritis in men. *Int J STD AIDS* 2014; **25**: 294-298.
31. Wetmore CM, Manhart LE, Lowens MS *et al.* *Ureaplasma urealyticum* is associated with nongonococcal urethritis among men with fewer lifetime sexual partners: A case-control study. *J Infect Dis* 2011; **204**: 1274-1282.
32. Yoshida T, Deguchi T, Meda S *et al.* Quantitative detection of *Ureaplasma parvum* (biovar 1) and *Ureaplasma urealyticum* (biovar 2) in urine specimens from men with and without urethritis by real-time polymerase chain reaction. *Sex Transm Dis* 2007; **34**: 416-419.
33. Huang C, Zhu HL, Xu KR, Wang SY, Fan LQ, Zhu WB. Mycoplasma and ureaplasma infection and male infertility: a systematic review and meta-analysis. *Andrology* 2015; **3**: 809-816.

34. Huang C, Long X, Jing S *et al.* *Ureaplasma urealyticum* and *Mycoplasma hominis* infections and semen quality in 19,098 infertile men in China. *World J Urol* 2016; **34**:1039-1044.
35. Ahmadi MH, Mirsalehian A, Sadighi Gilani MA, Bahador A, Talebi M. Asymptomatic infection with *Mycoplasma hominis* negatively affects semen parameters and leads to male infertility as confirmed by improved semen parameters after antibiotic treatment. *Urology* 2017; **100**:97-102.
36. Hay P, Patel S, Daniels D, Group FtBCE. UK National Guideline for the management of bacterial vaginosis 2012. Available at: <https://www.bashhguidelines.org/media/1041/bv-2012.pdf>
37. Zozaya M, Ferris MJ, Siren JD *et al.* Bacterial communities in penile skin, male urethra, and vaginas of heterosexual couples with and without bacterial vaginosis. *Microbiome* 2016; **4**: 16.
38. Mandar R, Punab M, Borovkova N *et al.* Complementary seminovaginal microbiome in couples. *Res Microbiol* 2015; **166**: 440-447.
39. van Oostrum N, De Sutter P, Meys J, Verstraelen H. Risks associated with bacterial vaginosis in infertility patients: a systematic review and meta-analysis. *Hum Reprod* 2013; **28**: 1809-1815.
40. Haahr T, Jensen JS, Thomsen L, Duus L, Rygaard K, Humaidan P. Abnormal vaginal microbiota may be associated with poor reproductive outcomes: a prospective study in IVF patients. *Hum Reprod* 2016; **31**: 795-803.
41. Abusarah EA, Awwad ZM, Charvalos E, Shehabi AA. Molecular detection of potential sexually transmitted pathogens in semen and urine specimens of infertile and fertile males. *Diagn Microbiol Infect Dis* 2013; **77**: 283-286.

42. Zeighami H, Peerayeh SN, Yazdi RS, Sorouri R. Prevalence of *Ureaplasma urealyticum* and *Ureaplasma parvum* in semen of infertile and healthy men. *Int J STD AIDS* 2009; **20**: 387-390.
43. Zhang L, Zhang KP, Liang CZ. *Ureaplasma urealyticum* in male genital tract: a hidden risk factor for male infertility. *Andrologia* 2016; **48**: 1077-1079.
44. Liu J, Wang Q, Ji X *et al.* Prevalence of *Ureaplasma urealyticum*, *Mycoplasma hominis*, *Chlamydia trachomatis* infections, and semen quality in infertile and fertile men in China. *Urology* 2014; **83**: 795-799.
45. Al-Sweih NA, Al-Fadli AH, Omu AE, Rotimi VO. Prevalence of *Chlamydia trachomatis*, *Mycoplasma hominis*, *Mycoplasma genitalium*, and *Ureaplasma urealyticum* infections and seminal quality in infertile and fertile men in Kuwait. *J Androl* 2012; **33**: 1323-1329.
46. Leli C, Mencacci A, Latino MA *et al.* Prevalence of cervical colonization by *Ureaplasma parvum*, *Ureaplasma urealyticum*, *Mycoplasma hominis* and *Mycoplasma genitalium* in childbearing age women by a commercially available multiplex real-time PCR: An Italian observational multicentre study. *J Microbiol Immunol Infect* 2017 Jun 28. pii: S1684-1182(17)30091-9.
47. Esen B, Gozalan A, Sevindi DF *et al.* *Ureaplasma urealyticum*: presence among sexually transmitted diseases. *Jpn J Infect Dis* 2017; **70**: 75-79.
48. Marovt M, Kese D, Kotar T *et al.* *Ureaplasma parvum* and *Ureaplasma urealyticum* detected with the same frequency among women with and without symptoms of urogenital tract infection. *Eur J Clin Microbiol Infect Dis* 2015; **34**: 1237-1245.
49. Lobao TN, Campos GB, Selis NN *et al.* *Ureaplasma urealyticum* and *U. parvum* in sexually active women attending public health clinics in Brazil. *Epidemiol Infect* 2017; **145**: 2341-2351.

50. Ahmadi MH, Mirsalehian A, Bahador A. Prevalence of urogenital Mycoplasmas in Iran and their effects on fertility potential: A systematic review and meta-analysis. *Iran J Public Health* 2016; **45**: 409-422.
51. Kyndel A, Elmer C, Kallman O, Altman D. Mycoplasmatataceae colonizations in women with urethral pain syndrome: A case-control study. *J Low Genit Tract Dis* 2016; **20**: 272-274.
52. Daxboeck F, Iro E, Tamussino K, Krause R, Assadian O, Wenisch C. Bacteremia with *Mycoplasma hominis* and *Ureaplasma urealyticum* in patients undergoing hysterectomy. *Eur J Clin Microbiol Infect Dis* 2003; **22**: 608-611.
53. Ruan Z, Yang T, Shi X, Kong Y, Xie X, Zhang J. Clonality and distribution of clinical *Ureaplasma* isolates recovered from male patients and infertile couples in China. *PLoS One* 2017; **12**: e0183947.
54. Ouzounova-Raykova VV, Markovska R, Mizgova G, Mitov IG. Detection of the sexually transmissible genital mycoplasmas by polymerase chain reaction in women. *Sex Health* 2011; **8**: 445-446.
55. Hammerschlag MR, Alpert S, Rosner I *et al.* Microbiology of the vagina in children: normal and potentially pathogenic organisms. *Pediatrics* 1978; **62**: 57-62.
56. Romero P, Muñoz M, Martínez MA *et al.* *Ureaplasmas* and mycoplasmas in vaginal samples from prepubertal girls and the reasons for gynecological consultation. *J Pediatr Adolesc Gynecol* 2014; 27:10-3.
57. McCormack WM, Almeida PC, Bailey PE, Grady EM, Lee YH. Sexual activity and vaginal colonization with genital mycoplasmas. *JAMA* 1972; **221**: 1375-1377.
58. Taylor-Robinson D, McCormack WM. The genital mycoplasmas (second of two parts). *N Engl J Med* 1980; **302**: 1063-1067.

59. Rosenstein IJ, Morgan DJ, Sheehan M, Lamont RF, Taylor-Robinson D. Bacterial vaginosis in pregnancy: distribution of bacterial species in different gram-stain categories of the vaginal flora. *J Med Microbiol* 1996; **45**: 120-126.
60. Cox C, Watt AP, McKenna JP, Coyle PV. *Mycoplasma hominis* and *Gardnerella vaginalis* display a significant synergistic relationship in bacterial vaginosis. *Eur J Clin Microbiol Infect Dis* 2016; **35**: 481-487.
61. Shipitsyna E, Roos A, Datcu R *et al.* Composition of the vaginal microbiota in women of reproductive age--sensitive and specific molecular diagnosis of bacterial vaginosis is possible? *PLoS One* 2013; **8**: e60670.
62. Ravel J, Gajer P, Abdo Z *et al.* Vaginal microbiome of reproductive-age women. *Proc Natl Acad Sci U S A* 2011; **108 Suppl 1**: 4680-4687.
63. Datcu R, Gesink D, Mulvad G *et al.* Vaginal microbiome in women from Greenland assessed by microscopy and quantitative PCR. *BMC Infect Dis* 2013; **13**: 480.
64. Austin MN, Beigi RH, Meyn LA, Hillier SL. Microbiologic response to treatment of bacterial vaginosis with topical clindamycin or metronidazole. *J Clin Microbiol* 2005; **43**: 4492-4497.
65. Pheifer TA, Forsyth PS, Durfee MA, Pollock HM, Holmes KK. Nonspecific vaginitis: role of *Haemophilus vaginalis* and treatment with metronidazole. *Engl J Med* 1978; **298**: 1429-1434.
66. Koutsky LA, Stamm WE, Brunham RC *et al.* Persistence of *Mycoplasma hominis* after therapy: importance of tetracycline resistance and of coexisting vaginal flora. *Sex Transm Dis* 1983; **10**: 374-381.
67. Lusk MJ, Konecny P, Naing ZW, Garden FL, Cumming RG, Rawlinson WD. *Mycoplasma genitalium* is associated with cervicitis and HIV infection in an urban Australian STI clinic population. *Sex Transm Infect* 2011; **87**: 107-109.

68. Lusk MJ, Garden FL, Rawlinson WD, Naing ZW, Cumming RG, Konecny P. Cervicitis aetiology and case definition: a study in Australian women attending sexually transmitted infection clinics. *Sex Transm Infect* 2016; **92**: 175-181.
69. Liu L, Cao G, Zhao Z, Zhao F, Huang Y. High bacterial loads of *Ureaplasma* may be associated with non-specific cervicitis. *Scand J Infect Dis* 2014; **46**: 637-641.
70. Stamm WE, Running K, Hale J, Holmes KK. Etiologic role of *Mycoplasma hominis* and *Ureaplasma urealyticum* in women with the acute urethral syndrome. *Sex Transm Dis* 1983; **10**: 318-322.
71. Lind K, Kristensen GB, Bollerup AC *et al.* Importance of *Mycoplasma hominis* in acute salpingitis assessed by culture and serological tests. *Genitourin Med* 1985; **61**: 185-189.
72. Michou IV, Constantoulakis P, Makarounis K, Georgoulas G, Kapetanios V, Tsilivakos V. Molecular investigation of menstrual tissue for the presence of *Chlamydia trachomatis*, *Ureaplasma urealyticum* and *Mycoplasma hominis* collected by women with a history of infertility. *J Obstet Gynaecol Res* 2014; **40**: 237-242.
73. Mårdh PA, Weström L. Tubal and cervical cultures in acute salpingitis with special reference to *Mycoplasma hominis* and T-strain mycoplasmas. *Br J Vener Dis* 1970; **46**: 179-186.
74. Taylor-Robinson D, Jensen JS, Svenstrup H, Stacey CM. Difficulties experienced in defining the microbial cause of pelvic inflammatory disease. *Int J STD AIDS* 2012; **23**:18-24.
75. Jurstrand M, Jensen JS, Magnuson A, Kamwendo F, Fredlund H. A serological study of the role of *Mycoplasma genitalium* in pelvic inflammatory disease and ectopic pregnancy. *Sex Transm Infect* 2007; **83**: 319-323.

76. Muin DA, Takes MT, Hosli I, Lapaire O. Severe pelvic abscess formation following caesarean section. *BMJ Case Rep* 2015 Apr 24;2015. pii: bcr2014208628.
77. Bailey EA, Solomon LR, Berry N *et al.* *Ureaplasma urealyticum* CAPD peritonitis following insertion of an intrauterine device: diagnosis by eubacterial polymerase chain reaction. *Perit Dial Int* 2002; **22**: 422-424.
78. Khosropour CM, Manhart LE, Gillespie CW *et al.* Efficacy of standard therapies against *Ureaplasma* species and persistence among men with non-gonococcal urethritis enrolled in a randomised controlled trial. *Sex Transm Infect* 2015; **91**: 308-313.
79. Taylor-Robinson D, Bébéar C. Antibiotic susceptibilities of mycoplasmas and treatment of mycoplasmal infections. *J Antimicrob Chemother* 1997; **40**: 622-630.
80. Beeton ML, Spiller OB. Antibiotic resistance among *Ureaplasma* spp. isolates: cause for concern? *J Antimicrob Chemother* 2017; **72**: 330-337.
81. Sfanos KS, Isaacs JT. The "infectious" nature of human prostate cancer: a cautionary note. *Oncotarget* 2011; **2**:281-283.
82. Barykova YA, Logunov DY, Shmarov MM *et al.* Association of *Mycoplasma hominis* infection with prostate cancer. *Oncotarget* 2011; **2**:289-297.
83. Rogers MB. Mycoplasma and cancer: in search of the link. *Oncotarget* 2011; **2**:271-273.
84. Lo SC, Tsai S. Mycoplasmas and human prostate cancer: an exciting but cautionary note. *Oncotarget* 2011; **2**:352-355.